

You can find the last version of these course materials at

<https://github.com/mbujosab/MatematicasII/tree/main/Eng>

- Marcos Bujosa. Copyright © 2008–2024

Algunos derechos reservados. Esta obra está bajo una licencia de Creative Commons Reconocimiento-CompartirIgual 4.0 Internacional. Para ver una copia de esta licencia, visite <http://creativecommons.org/licenses/by-sa/4.0/> o envie una carta a Creative Commons, 559 Nathan Abbott Way, Stanford, California 94305, USA.

Mathematics II

Marcos Bujosa

Universidad Complutense de Madrid

04/04/2024

1 / 23

1 / 23



1 Highlights of Lesson 13

Highlights of Lesson 13



- Determinant: $\det(\mathbf{A}) \equiv |\mathbf{A}|$ [det : $\mathbb{R}^{n \times n} \rightarrow \mathbb{R}$]
 - Volume vs determinant
 - Properties: [1](#), [2](#), [3](#)
- We will deduce properties: [4 – 9](#)

2 Area or volume



$$\text{1. } \text{Vol}\left(\mathbf{I}_{n \times n}\right) = 1.$$



$$\text{2. } \text{Vol}(\mathbf{A}) = \text{Vol}\left(\mathbf{A}_{[(\alpha)k+j]}\right) \text{ for } j \neq k.$$



$$\text{3. } |\alpha| \cdot \text{Vol}(\mathbf{A}) = |\alpha| \cdot \text{Vol}[\dots; \mathbf{A}_{[k]}; \dots] = \text{Vol}[\dots; \alpha \mathbf{A}_{[k]}; \dots]$$



3 Determinant: 3 properties that define the function

P-1 Determinant of identity matrices:

$$\det_{n \times n} \mathbf{I} = 1$$

P-2 Type I elemen. transf. do not change the determinant:

$$\det \mathbf{A} = \det \left(\mathbf{A}_{\tau_{[(\alpha)k+j]}} \right)$$

P-3 Multiplying a column by an scalar multiplies the det.

$\alpha \cdot \det \mathbf{A} = \det [\dots; \alpha \mathbf{A}_{|k}; \dots]$ for any $k \in \{1 : n\}$ and $\alpha \in \mathbb{R}$

Absolute value of $\det \mathbf{A} = \text{Vol } \mathbf{A}$

Example

Then, we know that in \mathbb{R}^3 :

$$\begin{vmatrix} a_1 & (b_1 + \alpha c_1) & c_1 \\ a_2 & (b_2 + \alpha c_2) & c_2 \\ a_3 & (b_3 + \alpha c_3) & c_3 \end{vmatrix} = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix};$$

$$\det [\mathbf{a}; (\mathbf{b} + \alpha \mathbf{c}); \mathbf{c}] = \det [\mathbf{a}; \mathbf{b}; \mathbf{c}];$$

and also

$$\begin{vmatrix} a_1 & \alpha b_1 & c_1 \\ a_2 & \alpha b_2 & c_2 \\ a_3 & \alpha b_3 & c_3 \end{vmatrix}; = \alpha \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix};$$

$$\det [\mathbf{a}; \alpha \mathbf{b}; \mathbf{c}] = \alpha \det [\mathbf{a}; \mathbf{b}; \mathbf{c}];$$

4 Determinant of a matrix with a zero column

P-4 Det. of a matrix \mathbf{A} with a zero column

If \mathbf{A} has a zero column $\mathbf{0}$, then

$$\det(\mathbf{A}) = 0$$

prove P-4

5 Elementary matrices

We already know

$$\det \left(\mathbf{A}_{\tau_{[(\alpha)k+j]}} \right) = |\mathbf{A}|; \quad \det \left(\mathbf{A}_{\tau_{[(\alpha)k]}} \right) = \alpha |\mathbf{A}|.$$

Determinant of elementary matrices

$$\det \left(\mathbf{I}_{\tau_{[(\alpha)k+j]}} \right) = 1 \quad \text{and} \quad \det \left(\mathbf{I}_{\tau_{[j]}} \right) = \alpha.$$

Hence, since $\mathbf{A}_\tau = \mathbf{A}(\mathbf{I}_\tau)$, then

$$|\mathbf{A}(\mathbf{I}_\tau)| = |\mathbf{A}| \cdot |\mathbf{I}_\tau| \tag{1}$$

where \mathbf{I}_τ is an elementary matrix

EXERCISE 1. Prove the following propositions

(a) $\det(\mathbf{A}_{\tau_1 \dots \tau_k}) = |\mathbf{A}| \cdot |\mathbf{I}_{\tau_1}| \cdots |\mathbf{I}_{\tau_k}|$.

(b) If \mathbf{B} is a full rank matrix, i.e., if $\mathbf{B} = \mathbf{I}_{\tau_1 \dots \tau_k}$, then $|\mathbf{B}| = |\mathbf{I}_{\tau_1}| \cdots |\mathbf{I}_{\tau_k}|$, and therefore $|\mathbf{B}| \neq 0$.

(c) If \mathbf{A} and \mathbf{B} have order n and \mathbf{B} is full rank, then

$$\det(\mathbf{AB}) = |\mathbf{A}| \cdot |\mathbf{B}| \quad (2)$$

7 Antisymmetric property

P-5 [Antisymmetric property]

Column exchange changes the sign of the determinant.

Proof.

Column exchange is a sequence of Type I transformation and just only one Type II transformation that multiplies a column by -1 \square

Therefore:

$$\begin{vmatrix} c_1 & b_1 & a_1 \\ c_2 & b_2 & a_2 \\ c_3 & b_3 & a_3 \end{vmatrix} = (-1) \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

6 Determinant after a sequence of elementary transformations

Example

a sequence $\tau_1 \dots \tau_k$ of Type I elementary transformations does not change the determinant.

$$|\mathbf{A}_{\tau_1 \dots \tau_k}| = |\mathbf{A}(\mathbf{I}_{\tau_1 \dots \tau_k})| = |\mathbf{A}| \cdot |\mathbf{I}_{\tau_1 \dots \tau_k}| = |\mathbf{A}| \cdot 1 = |\mathbf{A}|$$

Example

but a sequence of Type II can.

$$\begin{vmatrix} 2a & 3c \\ 2b & 3d \end{vmatrix} = ? \begin{vmatrix} a & c \\ b & d \end{vmatrix}$$

8 Singular matrices. Inverse of a matrix

P-6 If \mathbf{A} is singular then $|\mathbf{A}| = 0$.

$$\boxed{\text{P-7}} \quad \det(\mathbf{A}^{-1}) = (\det \mathbf{A})^{-1}.$$

Proof.

Let $\mathbf{A}_{\tau_1 \dots \tau_k} = \mathbf{R}$ be a reduced echelon form (and $\mathbf{E} = \mathbf{I}_{\tau_1 \dots \tau_k}$).

Since $\mathbf{AE} = \mathbf{R}$, then: $|\mathbf{A}| \cdot |\mathbf{E}| = |\mathbf{R}|$; with only two cases:

$$\begin{cases} \mathbf{A} \text{ singular } (\mathbf{R}_{|n} = 0) : & |\mathbf{A}| \cdot |\mathbf{E}| = 0 \Rightarrow |\mathbf{A}| = 0 \\ \mathbf{A} \text{ not singular } (\mathbf{R} = \mathbf{I}) : & |\mathbf{A}| \cdot |\mathbf{E}| = 1 \Rightarrow |\mathbf{E}| = |\mathbf{A}^{-1}| = (|\mathbf{A}|)^{-1} \end{cases}.$$

\square

Example

For $\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & 2 \end{bmatrix}$:

$$\begin{bmatrix} 1 & 2 \\ 2 & 2 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} \xrightarrow[\text{Type I}]{[(-2)\tau_1+2]} \begin{bmatrix} 1 & 0 \\ 2 & -2 \\ 1 & -2 \\ 0 & 1 \end{bmatrix} \xrightarrow[\text{Type II}]{[(-1)\tau_2]2} \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 1 & 1 \\ 0 & -1/2 \end{bmatrix} \xrightarrow[\text{Type I}]{[(-2)\tau_2+1]} \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ -1 & 1 \\ 1 & -1/2 \end{bmatrix}$$

So

$$|\mathbf{A}^{-1}| = \left| \mathbf{I}_{[(-2)\tau_1+2]} \right| \cdot \left| \mathbf{I}_{[(-1)\tau_2]2} \right| \cdot \left| \mathbf{I}_{[(-2)\tau_2+1]} \right| = 1 \cdot \frac{-1}{2} \cdot 1 = \frac{-1}{2};$$

that is $|\mathbf{A}| = -2.$

9 Determinant of a product**P-8****[Determinant of a product of matrices]**

$$\det(\mathbf{AB}) = \det(\mathbf{A}) \cdot \det(\mathbf{B}).$$

(3)

$$\begin{cases} \mathbf{B} \text{ singular, then so it is } \mathbf{AB} \Rightarrow \det(\mathbf{AB}) = 0 = \det(\mathbf{A}) \cdot \det(\mathbf{B}) \\ \mathbf{B} = \mathbf{I}_{\tau_1 \dots \tau_k} \Rightarrow \det(\mathbf{AB}) = \det(\mathbf{A}) \cdot \det(\mathbf{B}) \end{cases}$$

EXERCISE 2. [Transposed matrices]

- (a) What is the relation between the determinant of an elementary matrix \mathbf{I}_τ and the determinant of its transpose $\tau \mathbf{I}$?
 (b) Consider \mathbf{B} , a full rank matrix, proof that $|\mathbf{B}| = |\mathbf{B}^\top|$.

10 Determinant of a transpose**P-9****Determinant of a transpose**

$$|\mathbf{A}| = |\mathbf{A}^\top|.$$

Proof.

$$\begin{cases} \text{if } \mathbf{A} \text{ singular: } \mathbf{A}^\top \text{ singular } \Rightarrow \det \mathbf{A}^\top = \det \mathbf{A} = 0 \\ \text{if } \mathbf{A} \text{ NO singular: } \mathbf{A} = \mathbf{I}_{\tau_1 \dots \tau_k} \Rightarrow \det \mathbf{A}^\top = \det \mathbf{A} \end{cases}$$



Questions of the Lecture 13

(L-13) QUESTION 1. Complete the proofs of this lecture.

(L-13) QUESTION 2. Knowing that $|BC| = |B||C|$; prove that for any invertible matrix A (so $\det A \neq 0$)

$$\det(A^{-1}) = (\det(A))^{-1}.$$

(L-13) QUESTION 3. Consider $A_{3 \times 3}$ and $B_{3 \times 3}$ such that $\det(A) = 2$ and $\det(B) = -2$

- (a) (0.5pts) Compute the determinants of $A(B)^2$ and $(AB)^{-1}$
- (b) (0.5pts) Is it possible to compute the rank of $A + B$? and the rank of AB ?

(L-13) QUESTION 4. Use the Gauss-Jordan method to compute the determinant

$$(a) A_1 = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$(b) A_2 = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$$

$$(c) A_3 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

(L-13) QUESTION 5. The 3 by 3 matrix A reduces to the identity matrix I by the following three column operations (in order):

$\tau_{[(-4)1+2]}$: Subtract 4 times column 1 from column 2.

$\tau_{[(-3)1+3]}$: Subtract 3 times column 1 from column 3.

$\tau_{[(-1)3+2]}$: Subtract column 3 from column 2.

Find the determinant of A .

(L-13) QUESTION 6.

(a) Find the determinant of $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix}$ and $\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$

(b) Find the determinant of $\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ a & b & c & d \end{bmatrix}$ using Gauss-Jordan.

1 Highlights of Lesson 14

Highlights of Lesson 14

- Computing $|A|$ by gaussian elimination
- **P-10** — Multilinear property
- Expansion of $\det A$ in Cofactors (Laplace expansion).
- Application of determinants
 - Cramer's rule for solving linear equations
 - Computing the inverse of A

2 Extended matrix

Extended matrix of B : $\begin{bmatrix} B & 1 \end{bmatrix}$

1. Given τ : $\begin{bmatrix} B_\tau & 1 \end{bmatrix} = \begin{bmatrix} B & 1 \end{bmatrix}_\tau$.

2. Since $\begin{bmatrix} I & 1 \end{bmatrix}_\tau$ and I_τ same type Elem. Mat. \Rightarrow same det.

Applying 1. k times, and then 2.

$$\left| \begin{bmatrix} I_{\tau_1 \dots \tau_k} & 1 \end{bmatrix} \right| = \left| \begin{bmatrix} I & 1 \end{bmatrix}_{\tau_1 \dots \tau_k} \right| = \left| \begin{bmatrix} I & 1 \end{bmatrix}_{\tau_1} \dots \begin{bmatrix} I & 1 \end{bmatrix}_{\tau_k} \right| = \left| I_{\tau_1} \right| \dots \left| I_{\tau_k} \right| = \left| I_{\tau_1 \dots \tau_k} \right|.$$

If A is the extended matrix of B $\begin{cases} \text{If } B \text{ singular } & |B| = 0 = |A| \\ \text{If } B \text{ invertible } & |B| = |A| \end{cases}$

EXERCISE 7. [Triangular matrices]

- (a) Find the determinant of a full rank lower triangular matrix \mathbf{L}
- (b) Find the determinant of a triangular matrix with a zero entry in the main diagonal
- (c) Find the determinant of an upper triangular matrix \mathbf{U}

In addition
$$\begin{vmatrix} \mathbf{A} & \mathbf{0} \\ \mathbf{C} & \mathbf{B} \end{vmatrix}_{n \times m}^{m \times n} = |\mathbf{A}| \cdot |\mathbf{B}|.$$

Matrices of order 1, $\mathbf{A} = [a] :$

$$\begin{vmatrix} a & 0 \\ 0 & 1 \end{vmatrix} \Rightarrow |\mathbf{A}| = a.$$

Matrices of order 2:

$$\begin{vmatrix} a & b & 0 \\ c & d & 0 \\ 0 & 0 & 1 \end{vmatrix} \xrightarrow{\left[\begin{array}{l} (-\frac{b}{a})1+2 \\ (-\frac{c}{a})1+3 \end{array} \right]} \begin{vmatrix} a & 0 & 0 \\ c & d - \frac{bc}{a} & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

$$|\mathbf{A}| = ad - bc = a \det[d] - b \det[c].$$

Matrices of order 3:

$$\begin{vmatrix} a & b & c & 0 \\ d & e & f & 0 \\ g & h & i & 0 \\ 0 & 0 & 0 & 1 \end{vmatrix} \xrightarrow{\left[\begin{array}{l} (-\frac{b}{a})1+2 \\ (-\frac{c}{a})1+3 \\ (-\frac{e}{d})1+4 \end{array} \right]} \begin{vmatrix} a & 0 & 0 & 0 \\ d & e - \frac{bd}{a} & f - \frac{cd}{a} & 0 \\ g & h - \frac{bg}{a} & i - \frac{cg}{a} & 0 \\ 0 & 0 & 0 & 1 \end{vmatrix} \xrightarrow{\left[\begin{array}{l} (\frac{-af+cd}{ae-bd})2+3 \\ (\frac{-ah+de}{aj-be})2+4 \end{array} \right]}$$

$$\begin{vmatrix} a & 0 & 0 & 0 \\ d & e - \frac{bd}{a} & 0 & 0 \\ g & h - \frac{bg}{a} & \frac{ae-afh-bdi+bfg+cdh-ceg}{ae-bd} & 0 \\ 0 & 0 & 0 & 1 \end{vmatrix}$$

$$|\mathbf{A}| = \underbrace{aei - afh - bdi + bfg + cdh - ceg}_{(\text{Rule of Sarrus})} = a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix} - d \begin{vmatrix} e & f \\ i & j \end{vmatrix} + e \begin{vmatrix} f & g \\ k & l \end{vmatrix} - f \begin{vmatrix} g & h \\ m & o \end{vmatrix} + g \begin{vmatrix} h & i \\ p & o \end{vmatrix} - h \begin{vmatrix} i & j \\ m & n \end{vmatrix} + i \begin{vmatrix} j & k \\ n & o \end{vmatrix} - j \begin{vmatrix} k & l \\ m & n \end{vmatrix} + k \begin{vmatrix} l & m \\ n & o \end{vmatrix} - l \begin{vmatrix} m & n \end{vmatrix}$$

3 Computing by Gaussian elimination
Example

$$\mathbf{A} = \begin{bmatrix} 1 & 5 \\ 2 & 3 \end{bmatrix} : \begin{array}{c} \left[\begin{array}{cc|c} 1 & 5 & \\ 2 & 3 & \end{array} \right] \xrightarrow{\left[\begin{array}{l} (-5)1+2 \\ 1 \end{array} \right]} \left[\begin{array}{cc|c} 1 & 0 & \\ 2 & -7 & \end{array} \right] \end{array} \boxed{|\mathbf{A}| = -7}$$

Example

$$\begin{array}{c} \left[\begin{array}{ccc|c} 0 & 2 & 1 & 0 \\ 9 & 6 & 3 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right] \xrightarrow{\left[\begin{array}{l} (\frac{1}{2})3 \\ (-1)2+3 \\ (\frac{1}{2})4 \end{array} \right]} \left[\begin{array}{ccc|c} 0 & 2 & 0 & 0 \\ 9 & 6 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & \frac{1}{2} \end{array} \right] \xrightarrow{\left[\begin{array}{l} 1 \xrightarrow{=} 2 \\ (-1)4 \end{array} \right]} \left[\begin{array}{ccc|c} 2 & 0 & 0 & 0 \\ 6 & 9 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & -\frac{1}{2} \end{array} \right] \end{array}$$

$$\begin{vmatrix} 0 & 2 & 1 \\ 9 & 6 & 3 \\ 0 & 1 & 1 \end{vmatrix} = -9,$$

Matrices of order 4:

$$\begin{array}{c} \left[\begin{array}{cccc|c} a & b & c & d & 0 \\ e & f & g & h & 0 \\ i & j & k & l & 0 \\ m & n & o & p & 0 \end{array} \right] \xrightarrow{\left[\begin{array}{l} (-\frac{b}{a})1+2 \\ (-\frac{c}{a})1+3 \\ (-\frac{d}{a})1+4 \\ ((-\frac{f}{a})2+3) \\ ((-\frac{h}{a})2+4) \\ ((-\frac{b}{a})3+4) \end{array} \right]} \left[\begin{array}{cccc|c} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right] \end{array}$$

$$\begin{array}{c} \left[\begin{array}{cccc|c} a & 0 & 0 & 0 & 0 \\ e & f - \frac{be}{a} & 0 & 0 & 0 \\ i & j - \frac{bi}{a} & 0 & 0 & 0 \\ m & n - \frac{bm}{a} & 0 & 0 & 0 \end{array} \right] \xrightarrow{\left[\begin{array}{l} afk-agj-bek+beg+cej-cfi \\ af-o-agn-beo+bqm+cen-cfm \end{array} \right]} \left[\begin{array}{cccc|c} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right] \end{array}$$

$$\begin{array}{c} |\mathbf{A}| = \\ afkp - aflo - agjp + agln + ahjo - ahkn - bekp + belo + bgip - bglm - bhio + bhkm + cejp - celn - cfip + cflm + chin - chjm - dejo + dekn + dfio - dfkm - dgin + dgjm \\ = a \begin{vmatrix} f & g & h \\ j & k & l \\ n & o & p \end{vmatrix} - b \begin{vmatrix} e & g & h \\ i & k & l \\ m & o & p \end{vmatrix} + c \begin{vmatrix} e & f & h \\ i & j & l \\ m & n & p \end{vmatrix} - d \begin{vmatrix} e & f & g \\ i & j & k \\ m & n & o \end{vmatrix} \end{array}$$

4 Multilinear property

P-10 Multilinear property

$$\det [\dots; (\beta \mathbf{b} + \psi \mathbf{c}); \dots] = \beta \det [\dots; \mathbf{b}; \dots] + \psi \det [\dots; \mathbf{c}; \dots]$$

Example

Then, in the 2 dimensional case \mathbb{R}^2

$$\begin{vmatrix} a+\alpha & c \\ b+\beta & d \end{vmatrix} = \begin{vmatrix} a & c \\ b & d \end{vmatrix} + \begin{vmatrix} \alpha & c \\ \beta & d \end{vmatrix};$$

therefore

$$\begin{vmatrix} a & c \\ b & d \end{vmatrix} = \begin{vmatrix} a & c \\ 0 & d \end{vmatrix} + \begin{vmatrix} c \\ d \end{vmatrix}.$$

Example

For $\mathbf{A} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$, we have

$${}^1\mathbf{A}^{\hat{2}} = \begin{bmatrix} 4 & 6 \\ 7 & 9 \end{bmatrix}, \quad {}^3\mathbf{A}^{\hat{3}} = \begin{bmatrix} 1 & 2 \\ 4 & 5 \end{bmatrix}$$

hence

$$\text{cof}_{12}(\mathbf{A}) = (-1)^{1+2} \det({}^1\mathbf{A}^{\hat{2}}) = (-1) \begin{vmatrix} 4 & 6 \\ 7 & 9 \end{vmatrix}.$$

and

$$\text{cof}_{33}(\mathbf{A}) = (-1)^{3+3} \det({}^3\mathbf{A}^{\hat{3}}) = \begin{vmatrix} 1 & 2 \\ 4 & 5 \end{vmatrix}.$$

5 minors and cofactors

Definition minors and cofactors

We denote a submatrix of \mathbf{A} obtained by deleting row i and column j of \mathbf{A} by

$${}^i\mathbf{A}^{\hat{j}};$$

Its determinant is called the minor of a_{ij} . And

$$\text{cof}_{ij}(\mathbf{A}) = (-1)^{i+j} \det({}^i\mathbf{A}^{\hat{j}})$$

is called the cofactor of a_{ij} .

6 Expansion by cofactors

Theorem [Laplace expansion]

For \mathbf{A} n by n , $\det(\mathbf{A})$ may be computed as the sum of the products of the elements of any column (row) of \mathbf{A} by their cofactors:

$$\det(\mathbf{A}) = \sum_{i=1}^n a_{ij} \text{cof}_{ij}(\mathbf{A}), \quad \text{the expansion by the } j\text{th column}$$

or

$$\det(\mathbf{A}) = \sum_{j=1}^n a_{ij} \text{cof}_{ij}(\mathbf{A}), \quad \text{the expansion by the } i\text{th row}$$

7 Cramer's Rule

$$\mathbf{A}\mathbf{x} = \mathbf{b}; \quad |\mathbf{A}| \neq 0 \quad \text{then}$$

$$\mathbf{b} = (\mathbf{A}_{|1})x_1 + \cdots + (\mathbf{A}_{|j})\overset{\text{pos. } j}{\overbrace{\mathbf{x}_j}} + \cdots + (\mathbf{A}_{|n})x_n.$$

EXERCISE 8. Compute $\det \mathbf{A} = \begin{vmatrix} 2 & 0 & 3 & 2 \\ 5 & 1 & 2 & 4 \\ 3 & 0 & 1 & 2 \\ 5 & 3 & 2 & 1 \end{vmatrix}$

$$\det \left[\mathbf{A}_{|1}; \dots, \overset{\text{pos. } j}{\overbrace{\mathbf{b}}}; \dots, \mathbf{A}_{|n} \right] = \mathbf{x}_j \cdot \det(\mathbf{A}).$$

$$x_j = \frac{\det \left[\mathbf{A}_{|1}; \dots, \overset{\text{pos. } j}{\overbrace{\mathbf{b}}}; \dots, \mathbf{A}_{|n} \right]}{\det(\mathbf{A})}.$$

Computational issues when $\det \mathbf{A} \approx 0$ (tiny angle between vectors)

8 The inverse of a matrix

$$[\text{Adj}(\mathbf{A})] \cdot \mathbf{A} =$$

$$\begin{bmatrix} \text{cof}_{11}(\mathbf{A}) & \text{cof}_{21}(\mathbf{A}) & \cdots & \text{cof}_{n1}(\mathbf{A}) \\ \text{cof}_{12}(\mathbf{A}) & \text{cof}_{22}(\mathbf{A}) & \cdots & \text{cof}_{n2}(\mathbf{A}) \\ \vdots & \vdots & \ddots & \vdots \\ \text{cof}_{1n}(\mathbf{A}) & \text{cof}_{2n}(\mathbf{A}) & \cdots & \text{cof}_{nn}(\mathbf{A}) \end{bmatrix} \underbrace{\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}}_{\mathbf{A}}$$

Questions of the Lecture 14

(L-14) QUESTION 1. Complete the proofs of the exercises of this lecture.

(L-14) QUESTION 2. Consider $\mathbf{A} = [\mathbf{A}_{|1}; \mathbf{A}_{|2}; \mathbf{A}_{|3}]$ with $\det \mathbf{A} = 2$.

- (a) What are $\det(2\mathbf{A})$ and $\det \mathbf{A}^{-1}$?
- (b) What is $\det [(3\mathbf{A}_{|1} + 2\mathbf{A}_{|2}); \mathbf{A}_{|3}; \mathbf{A}_{|2}]$?

(L-14) QUESTION 3. The determinant of the 1000 by 1000 matrix \mathbf{A} is 12. What is the determinant of $-\mathbf{A}^T$? (Careful: No credit for the wrong sign.)
(MIT Course 18.06 Quiz 2, Fall, 2008)

(L-14) QUESTION 4. Consider the squared matrix \mathbf{A} . True or false? (to receive full credit you must explain your answer in a clear and concise way)
 $|\mathbf{A}\mathbf{A}^T| = |\mathbf{A}|^2$.

(L-14) QUESTION 5. We have a 3×3 matrix $\mathbf{A} = \begin{bmatrix} a & 1 & 2 \\ b & 3 & 4 \\ c & 5 & 6 \end{bmatrix}$ with $\det \mathbf{A} = 3$.

Compute the determinant of the following matrices:

(a) (0.5 pts) $\begin{bmatrix} a-2 & 1 & 2 \\ b-4 & 3 & 4 \\ c-6 & 5 & 6 \end{bmatrix}$

(b) (0.5 pts) $\begin{bmatrix} 7a & 7 & 14 \\ b & 3 & 4 \\ c & 5 & 6 \end{bmatrix}$

(c) (1 pts) $(2\mathbf{A})^{-1}\mathbf{A}^T$
(d) (0.5 pts) $\begin{bmatrix} a-2 & 1 & 2 \\ b & 3 & 4 \\ c & 5 & 6 \end{bmatrix}$

(L-14) QUESTION 6.

(a) Escalone la matriz $\mathbf{A} = \begin{bmatrix} 2 & 3 & 1 \\ 4 & 5 & 2 \\ 4 & 6 & 0 \end{bmatrix}$.

(b) ¿Es \mathbf{A} invertible?

(c) En caso afirmativo calcule $|\mathbf{A}^{-1}|$; en caso contrario calcule $|\mathbf{A}|$

(d) La matriz \mathbf{C} es igual al producto de \mathbf{A} con la *traspuesta* de la matriz \mathbf{B} , es decir

$$\mathbf{C} = \mathbf{AB}^T \quad \text{donde} \quad \mathbf{B} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 0 & 2 & 3 \end{bmatrix}$$

¿Cuánto vale el determinante de \mathbf{C} ? ¿Es \mathbf{C} invertible?

(L-14) QUESTION 7. What is the determinant of the following matrices using Laplace expansions.

(a) $\begin{bmatrix} 1 & 2 \\ -4 & 3 \end{bmatrix}$

(b) $\begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & 2 \\ 2 & 1 & -2 \end{bmatrix}$

(c) $\begin{bmatrix} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 2 \\ 2 & 0 & 1 & -2 \end{bmatrix}$

(L-14) QUESTION 8. Compute the following determinant using Laplace expansions:

$$\begin{vmatrix} 0 & 0 & 0 & 3 & 0 \\ -2 & 0 & 0 & 2 & 0 \\ 8 & -1 & 0 & -7 & 2 \\ -1 & 2 & 2 & 3 & 2 \\ 2 & 2 & 3 & 6 & 4 \end{vmatrix}$$

(L-14) QUESTION 9. Compute $\det \mathbf{A} =$

$$\begin{vmatrix} 2 & 2 & 0 & 0 \\ 5 & 5 & 0 & 0 \\ 3 & 0 & 1 & 0 \\ 5 & 0 & 0 & 1 \end{vmatrix}$$

(L-14) QUESTION 10. Compute the value of $\det \mathbf{A}$ using Laplace expansion

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 \\ 0 & 2 & 2 & \cdots & 2 \\ 0 & 0 & 3 & \cdots & 3 \\ \vdots & & \ddots & & \\ 0 & 0 & 0 & \cdots & n \end{bmatrix}$$

(L-14) QUESTION 11. Consider a n by n matrix \mathbf{A}_n full of 3s in its diagonal, and twos just below the diagonal, and another 2 at the position $(1, n)$; for example, for $n = 4$:

$$\mathbf{A}_4 = \begin{bmatrix} 3 & 0 & 0 & 2 \\ 2 & 3 & 0 & 0 \\ 0 & 2 & 3 & 0 \\ 0 & 0 & 2 & 3 \end{bmatrix}.$$

- (a) Find, using the cofactors of the first row, the determinant of \mathbf{A}_4 .
(b) Find the determinant of \mathbf{A}_n for $n > 4$.

(L-14) QUESTION 12. Consider the following block matrix

$$\mathbf{A} = \begin{bmatrix} \mathbf{B} & \mathbf{0} \\ \mathbf{0} & \mathbf{C} \end{bmatrix}$$

Prove $|\mathbf{A}| = |\mathbf{B}||\mathbf{C}|$.

Hint

$$\begin{bmatrix} \mathbf{B} & \mathbf{0} \\ \mathbf{0} & \mathbf{C} \end{bmatrix} = \begin{bmatrix} \mathbf{B} & \mathbf{0} \\ \mathbf{0} & \mathbf{I} \end{bmatrix} \begin{bmatrix} \mathbf{I} & \mathbf{0} \\ \mathbf{0} & \mathbf{C} \end{bmatrix}$$

(L-14) QUESTION 13. Solve the following linear systems using Cramer's Rule

(a) $\begin{cases} 2x + 5y = 1 \\ x + 4y = 2 \end{cases}$

(b) $\begin{cases} 2x + y = 1 \\ x + 2y + z = 0 \\ y + 2z = 0 \end{cases}$

(exercise 13 from section 4.4 of Strang (2006))

(L-14) **QUESTION 14.** Find the inverse of the following matrices using the *adjoint matrix*

(a) $\mathbf{A} = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 3 & 0 \\ 0 & 4 & 1 \end{bmatrix}$

(b) $\mathbf{B} = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$

(exercise 18 from section 4.4 of Strang (2006))

(L-14) **QUESTION 15.** Consider the matrices

$$\mathbf{A} = \begin{bmatrix} 1 & 4 & 2 & 3 \\ 2 & 3 & 3 & 7 \\ 0 & 1 & 0 & 3 \\ 0 & 2 & 0 & a \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{bmatrix}; \text{ and the vector } \mathbf{b} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}.$$

(a) (0.5pts) For which values of a the matrix \mathbf{A} is invertible?

(b) (1pts) Consider $a = 5$. Using the Cramer's rule, compute the fourth coordinate x_4 of \mathbf{x} for linear system $\mathbf{Ax} = \mathbf{b}$.

(c) (1pts) Compute \mathbf{B}^{-1} . Use the matrix \mathbf{B}^{-1} to solve $\mathbf{Bx} = \mathbf{b}$.

Strang, G. (2006). *Linear algebra and its applications*. Thomson Learning, Inc., fourth ed. ISBN 0-03-010567-6.