

Mathematics II

Marcos Bujosa

Universidad Complutense de Madrid

04/04/2024

1 / 15

L-19

You can find the last version of these course materials at

<https://github.com/mbujosab/MatematicasII/tree/main/Eng>

– Marcos Bujosa. Copyright © 2008–2024
Algunos derechos reservados. Esta obra está bajo una licencia de Creative Commons Reconocimiento-CompartirIgual 4.0 Internacional. Para ver una copia de esta licencia, visite <http://creativecommons.org/licenses/by-sa/4.0/> o envíe una carta a Creative Commons, 559 Nathan Abbott Way, Stanford, California 94305, USA.

1 / 15

L-19

1 Highlights of Lesson 19

Highlights of Lesson 19

- Mean
- Standard deviation and variance
- Ordinary Least Squares (OLS)

2 / 15

L-19

2 Restriction in statistics and probability

Norm of constant vector “one” is 1

This fails using the dot product in \mathbb{R}^m ($m > 1$)

$$\|\mathbf{1}\|^2 = \langle \mathbf{1} | \mathbf{1} \rangle = \mathbf{1} \cdot \mathbf{1} = \sum_{i=1}^m 1 = m.$$

New scalar product in \mathbb{R}^m for statistics

$$\langle \mathbf{x} | \mathbf{y} \rangle_s = \frac{1}{m} (\mathbf{x} \cdot \mathbf{y})$$

(so: $\|\mathbf{1}\|^2 = \frac{1}{m} (\mathbf{1} \cdot \mathbf{1}) = 1$)

3 / 15

3 Mean

The mean $\mu_{\mathbf{y}}$ is the scalar product of \mathbf{y} and $\mathbf{1}$

$$\mu_{\mathbf{y}} = \frac{1}{m}(\mathbf{1} \cdot \mathbf{y}), \quad \text{so, } \mu_{\mathbf{y}} = \frac{1}{m} \sum_{i=1}^m y_i$$

The mean $\mu_{\mathbf{y}}$ is the *value* by which to multiply $\mathbf{1}$ to get the orthogonal projection of \mathbf{y} onto $\mathcal{L}([\mathbf{1};])$

$\bar{\mathbf{y}}$: projection of $\mathbf{y} \in \mathbb{R}^m$ onto the line $\mathcal{L}([\mathbf{1};]) \subset \mathbb{R}^m$


$$\bar{\mathbf{y}} = \mathbf{1}\hat{a} \quad \text{and} \quad (\mathbf{y} - \bar{\mathbf{y}}) \perp \mathbf{1} \Rightarrow \frac{1}{m}(\mathbf{y} - \bar{\mathbf{y}}) \cdot \mathbf{1} = 0$$

$$\frac{1}{m}(\mathbf{y} - \mathbf{1}\hat{a}) \cdot \mathbf{1} = 0 \Leftrightarrow \frac{1}{m}(\mathbf{y} \cdot \mathbf{1}) - \frac{1}{m}(\mathbf{1} \cdot \mathbf{1})\hat{a} = 0;$$

Therefore


$$\hat{a} = \frac{1}{m}(\mathbf{y} \cdot \mathbf{1}) = \mu_{\mathbf{y}}$$

4 Mean




5 Standard deviation

$$\sigma_{\mathbf{y}} = \|\mathbf{y} - \bar{\mathbf{y}}\|.$$



6 Constant Vectors and Zero Mean Vectors


Adding a constant vector $a\mathbf{1}$ to \mathbf{y} does not change the standard deviation.



$$\sigma_{\mathbf{z}} = 0 \Leftrightarrow \mathbf{z} = a\mathbf{1}; \quad \mu_{\mathbf{z}} = 0 \Leftrightarrow \mathbf{z} \perp \mathbf{1}$$

7 Variance and the Pythagorean theorem

$$\sigma_y^2 = \|\mathbf{y} - \bar{\mathbf{y}}\|^2 = \frac{1}{m}(\mathbf{y} - \bar{\mathbf{y}}) \cdot (\mathbf{y} - \bar{\mathbf{y}}) = \frac{1}{m} \sum_i (y_i - \mu_y)^2.$$




$$\sigma_y^2 = \|\mathbf{y} - \bar{\mathbf{y}}\|^2 = \|\mathbf{y}\|^2 - \|\bar{\mathbf{y}}\|^2 = \frac{1}{m}(\mathbf{y} \cdot \mathbf{y}) - \mu_y^2 = \frac{\sum_i y_i^2}{m} - \mu_y^2.$$

8 / 15

8 Covariance and correlation

$$\sigma_{xy} = \frac{1}{m}(\mathbf{x} - \mu_x) \cdot (\mathbf{y} - \bar{\mathbf{y}});$$



$$\rho_{xy} = \frac{\frac{1}{m}(\mathbf{x} - \mu_x) \cdot (\mathbf{y} - \bar{\mathbf{y}})}{\|\mathbf{x} - \mu_x\| \cdot \|\mathbf{y} - \bar{\mathbf{y}}\|} = \frac{\sigma_{xy}}{\sqrt{\sigma_x \sigma_y}} = \cos(\theta).$$

9 / 15

9 Ordinary Least Squares (OLS)

Let \mathbf{X} such that $\mathcal{L}([1;]) \subset \mathcal{C}(\mathbf{X})$.

$\hat{\mathbf{y}}$ is the orthogonal projection of $\mathbf{y} \in \mathbb{R}^m$ onto $\mathcal{C}(\mathbf{X})$

$$\hat{\mathbf{y}} = \mathbf{X}\hat{\beta} \quad \text{and} \quad (\mathbf{y} - \hat{\mathbf{y}}) \perp \mathcal{C}(\mathbf{X}) \Rightarrow \frac{1}{m}\mathbf{X}^\top(\mathbf{y} - \hat{\mathbf{y}}) = \mathbf{0}$$

$$\frac{1}{m}\mathbf{X}^\top(\mathbf{y} - \mathbf{X}\hat{\beta}) = \mathbf{0} \iff \frac{1}{m}\mathbf{X}^\top\mathbf{y} - \frac{1}{m}\mathbf{X}^\top\mathbf{X}\hat{\beta} = \mathbf{0}.$$


Therefore

$$\left(\frac{1}{m}\mathbf{X}^\top\mathbf{X}\right)\hat{\beta} = \frac{1}{m}\mathbf{X}^\top\mathbf{y}.$$

10 / 15

10 Ordinary Least Squares (OLS)

If $\mathbf{X} = [1; x;]$ has rank 2.




$$\left(\frac{1}{m}\mathbf{X}^\top\mathbf{X}\right) \begin{pmatrix} \hat{a} \\ \hat{b} \end{pmatrix} = \frac{1}{m}\mathbf{X}^\top\mathbf{y}.$$

11 / 15

11 Application: Least Squares (Fitting by a line)

"looking for the best fitting line $\hat{y} = \hat{a} + \hat{b}x$ "

Points (x, y) : (1, 1); (2, 2); (3, 2)



$$\begin{cases} a + 1b = 1 \\ a + 2b = 2 \\ a + 3b = 2 \end{cases} \rightarrow \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \end{bmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix} \quad (\mathbf{X}\boldsymbol{\beta} = \mathbf{y} \text{ No solution})$$

12/15

12 Application: Least Squares (Fitting by a line)


$$\mathbf{X}\boldsymbol{\beta} = \mathbf{y} \quad (\text{No solution}) \rightarrow \left(\frac{1}{m}\mathbf{X}^T\mathbf{X}\right)\hat{\boldsymbol{\beta}} = \frac{1}{m}\mathbf{X}^T\mathbf{y} \rightarrow \hat{\mathbf{y}} = \mathbf{X}\hat{\boldsymbol{\beta}}.$$

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \end{bmatrix} \begin{pmatrix} \hat{a} \\ \hat{b} \end{pmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \end{bmatrix} \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$$

$$\begin{bmatrix} 3 & 6 \\ 6 & 14 \end{bmatrix} \begin{pmatrix} \hat{a} \\ \hat{b} \end{pmatrix} = \begin{pmatrix} 5 \\ 11 \end{pmatrix} \Rightarrow \hat{a} = \frac{2}{3}; \quad \hat{b} = \frac{1}{2}.$$

Best solution: $\frac{2}{3} + \frac{1}{2}x$

13/15

13 Application: Least Squares (Fitting by a line)

$$\hat{\mathbf{y}} = \mathbf{X}\hat{\boldsymbol{\beta}} = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \end{bmatrix} \begin{pmatrix} \hat{a} \\ \hat{b} \end{pmatrix} \rightarrow \hat{\mathbf{y}} = \begin{pmatrix} 7/6 \\ 10/6 \\ 13/6 \end{pmatrix} \rightarrow \hat{\mathbf{e}} = \begin{pmatrix} -1/6 \\ 2/6 \\ -1/6 \end{pmatrix}$$

$$\mathbf{y} = \hat{\mathbf{y}} + \hat{\mathbf{e}} \quad \text{and} \quad \begin{cases} \hat{\mathbf{e}} \cdot \hat{\mathbf{y}} = 0 \\ \hat{\mathbf{e}}\mathbf{X} = \mathbf{0} \end{cases}$$


14/15

14 Application: Least Squares (Fitting by a line)

Selling price and living area of single family homes in University City community of San Diego, in 1990.

price = Sale price is in thousands of dollars

sqft = Square feet of living area (Ramanathan, 2002, pp. 78)



15/15

Questions of the Lecture 19

(L-19) QUESTION 1. With the measurements $\mathbf{y} = (0, 8, 8, 20,)$ at $\mathbf{x} = (0, 1, 3, 4,)$,

- Set up and solve the normal equations $\mathbf{A}^T \mathbf{A} \hat{\boldsymbol{\beta}} = \mathbf{A}^T \mathbf{y}$.
- For the best straight line, find its four fits p_i and four errors e_i .
- What is the value of the square of the norm of the error vector $\|\mathbf{e}\|^2 = e_1^2 + e_2^2 + e_3^2 + e_4^2$?
- Draw the regression line
- Change the measurements to $\mathbf{p} = (1, 5, 13, 17,)$ write down the four equations $\mathbf{A}\boldsymbol{\beta} = \mathbf{p}$. Find an exact solution to $\mathbf{A}\boldsymbol{\beta} = \mathbf{p}$
- Check that $\mathbf{e} = \mathbf{y} - \mathbf{p} = (-1, 3, -5, 3,)$ is perpendicular to both columns of the same matrix \mathbf{A} .
- What is the shortest distance $\|\mathbf{e}\|$ from \mathbf{y} to the column space of \mathbf{A} ?

(Strang, 2003, exercise 1–3 from section 4.3.)

(L-19) QUESTION 2.

- Write down three equations $y = \alpha + \beta x$ given the data: $y = 7$ at $x = -1$, $y = 7$ at $x = 1$, and $y = 21$ at $x = 2$. Find the least squares solution $\hat{\boldsymbol{\beta}} = (\hat{\alpha}, \hat{\beta})$ and draw the closest line.
- Find the projection $\mathbf{p} = \mathbf{A}\hat{\boldsymbol{\beta}}$. This gives the three heights of the closest line. Show that the error vector is $\mathbf{e} = (2, -6, 4,)$. Why is $\mathbf{P}\mathbf{e} = \mathbf{0}$?

(L-19) QUESTION 3. Our measurements at times $t = 1, 2, 3$ are $b = 1, 4$, and b_3 . We want to fit those points by the nearest line $C + Dt$, using least squares.

- Which value for b_3 will put the three measurements on a straight line? Which line is it? Will least squares choose that line if the third measurement is $b_3 = 9$? (Yes or no).
- What is the linear system $\mathbf{A}\mathbf{x} = \mathbf{b}$ that would be solved exactly for $\mathbf{x} = (C, D)$ if the three points do lie on a line? Compute the projection matrix \mathbf{P} onto the column space of \mathbf{A} .
- What is the rank of that projection matrix \mathbf{P} ? How is the column space of \mathbf{P} related to the column space of \mathbf{A} ? (You can answer with or without the entries of \mathbf{P} computed in (b).)
- Suppose $b_3 = 1$. Write down the equation for the best least squares solution $\hat{\mathbf{x}}$, and show that the best straight line is horizontal.

Ramanathan, R. (2002). *Introductory Econometrics with applications*. South-Western, Mason, Ohio, fifth ed. ISBN 0-03-034186-8.

Strang, G. (2003). *Introduction to Linear Algebra*. Wellesley-Cambridge Press, Wellesley, Massachusetts. USA, third ed. ISBN 0-9614088-9-8.